
Abstract:

Emergency medicine practitioners often see young patients who are treated for injuries sustained during a violent encounter, most often with a peer from the same neighborhood. In addition, many more of the children and adolescents that we see are affected by the violence that surrounds them in their homes, neighborhood, and schools. This article reviews the prevalence and impact of interpersonal violence on our young patients, offers a suggested management approach to assault-injured children and adolescents who visit the emergency department, and reviews multidisciplinary outpatient programs for which the emergency department practitioners can advocate within their medical and social services systems.

Keywords:

violence; adolescence; assault; trauma; emergency medicine

*Emergency Department, The Children's Hospital of Philadelphia, Philadelphia, PA; †The Perelman School of Medicine at The University of Pennsylvania, Philadelphia, PA; ‡Greene Consulting, Montclair, NJ. Reprint requests and correspondence: Joel A. Fein, MD, MPH, Emergency Department Offices CTRB, 9th Floor, The Children's Hospital of Philadelphia, 3501 Civic Center Blvd, Philadelphia, PA 19104. fein@email.chop.edu (J.A. Fein) URL: <http://www.greeneconsultingnj.com> (M.B. Greene)

1522-8401/\$ - see front matter
© 2013 Elsevier Inc. All rights reserved.

The Assault-Injured Youth and the Emergency Medical System: What Can We Do?

Joel A. Fein, MD, MPH*†,
Cynthia J. Mollen, MD, MSCE*†,
Michael B. Greene, PhD‡

Interpersonal violence occurs between 2 or more noncare-taker individuals in which at least one individual intended to harm the other. These altercations frequently occur in the school, schoolyard, or street. Interpersonal violence differs from family violence, such as child abuse and domestic violence, in which an individual has a significant power or caretaking responsibilities over another within the relationship. Although most health care systems have protocols for the management of child abuse and some for domestic violence, there is no mandated reporting system or accepted psychosocial protocol for patients who are injured through interpersonal violence.

The 2 most widely understood facts about youth violence are as follows: (1) violence victimization and perpetration peak during the adolescent and young adult years, and (2) a very small percentage of adolescents perpetrate the most serious forms of violence, and correspondingly, a very small percentage of adolescents require medical attention as a consequence of violent

victimization.^{1,2} Less well known is the fact that there is significant overlap among adolescent victims and perpetrators; victims of violence are more likely to have histories and subsequent likelihoods of violence perpetration and vice versa.³ Being a victim of physical assault increases the risk of subsequent violent offending by up to 350%.⁴ Indeed, these individuals tend to have a common set of risk factors and engage in similar lifestyle activities in high-crime areas.⁵ The phenomenon of the same individuals appearing repeatedly in the same hospital has led to much frustration among emergency department (ED) physicians⁶ and has prompted the American Academy of Pediatrics to issue a model protocol to address the needs of adolescent assault victims.⁷ In that same year, the Society of Academic Emergency Medicine issued its own report and recommendations regarding the role that the emergency physician can play in reducing subsequent violence among assaulted victims treated in the ED.⁸

Clearly, serious violent injury provides a tragic and potentially teachable moment in an adolescent's life.⁹ Moreover, there is growing evidence and consensus that much can be done in the hospital setting to reduce the rate of injury recurrence and subsequent retaliatory violence.¹⁰

A NATIONAL PROBLEM

Interpersonal violence remains a major issue in American society. Homicide is the second leading cause of death for all Americans aged 15 to 24 years, accounting for almost 4700 deaths in this age group in 2010, a statistic that is unchanged in more than a decade.¹¹ Homicide rates do not tell the entire story, however; in 2011, almost 800 000 youth aged 15 to 24 years were cared for in an ED for injuries caused by violence, and 11% of these patients were hospitalized.¹² In urban communities, interpersonal intentional injuries account for 25% of all youth injuries, 45% of hospitalizations, and 85% of injury deaths.¹³ However, children from all settings are vulnerable; one study found that 89% of students in a suburban school knew someone who had been robbed, beaten, stabbed, shot, or murdered, and 57% had witnessed such an event. In a comparative urban school, 96% of students knew the victim of a violent crime, and 88% had witnessed an attack.¹⁴ Similarly, in a study of multiple towns and cities in Connecticut, although a higher proportion of poor, urban children witnessed violence, those in non-poor, suburban communities were not immune.¹⁵ Johnson and colleagues¹⁶ report that rural teens were as or more likely than urban and suburban

teens to display violent behavior or experience victimization.

Importantly, even homicide rates combined with hospital visits do not paint a complete picture of violence-related morbidity. Recently, more subtle effects of "indirect" exposures to violence have been identified. Adolescents, especially girls, who witness violence are more likely to experience symptoms of posttraumatic stress disorder than are adolescents who do not witness violent events.¹⁷

ASSESSMENT OF THE ASSAULT-INJURED YOUTH

Given the significant impact of interpersonal violence, we need to consider our assessment of assault-injured youth. Similar to how we assess patients with asthma, diabetes, or other illnesses for their risk of returning in similar or worse condition, we can try to assess how likely the youth is to return with another violent injury or subsequently injure another individual, frequently motivated by retaliation and a norm of retribution.¹⁸ This assessment can be divided into 3 components: a brief screening for immediate safety risk, a screening instrument to identify longer-term psychosocial risk, and a more thorough assessment of problem areas identified by the screening instrument. Based on the results from such assessments, risk and protective factors can be identified and a posthospital release plan established.

A systematic and sensitive approach to questioning adolescents removes blame and judgment, discusses confidentiality and reporting requirements, and engenders the adolescent's trust that the ED team is interested in his or her safety and well-being. The entire ED staff should be well attuned to the complexities that may lead to a single violent injury.⁶ As mentioned, it is not useful to apply the terms "victim" and "perpetrator" because, often, the "victim" that presents to the ED may have instigated the fight that he or she subsequently "lost." Receptive and positive attitudes are key: adolescents do not generally view the ED as the appropriate place to be counseled about violence.¹⁹ They are remarkably attuned to nonverbal and verbal cues about the feelings of the adults around them, and it is important to convey to them that the downward path into ever increasing and repeated violence is not inevitable. To write these patients off as hopeless or forever caught up in the mire of violence is akin to a self-fulfilling prophecy.^{20,21} Although they, like all of us, are responsible for their actions, multiple factors contribute to violence, some of which are out of their control.

Therefore, a “trauma-informed” approach to the violently injured patient is necessary to avoid increasing the impact of the event on their presenting complaints, their reactions to treatment interventions, and their adherence to discharge recommendations.²² This approach takes into consideration how community and institutional stressors such as poverty and racism and individual-level stressors such as poor nutrition, family dysfunction, and family violence influence how an adolescent copes with a threatening situation. The response to stress, driven more by the limbic system than by the cortex, needs to be considered when discussing how we counsel an assault-injured youth on how to avoid retaliation or risky situations.²³ For example, asking “what happened to you?” rather than “what’s wrong with you?” or “what did you do?” avoids the perception that they are being judged and creates a dialogue rather than an interrogation. The patient can be retraumatized if not allowed some control over the depth and direction of the conversation about event.²⁴

Trauma-informed emergency providers have also learned to recognize acute stress reactions, which can be additive to these conditions and complicate this picture.^{25,26} These include symptoms of hyperarousal, reexperiencing the event, dissociation (feeling like you were not really “there”), and avoidance.

Brief Screen for Immediate Safety Risk

The Immediate Safety Screen for Violently Injured Youth outlines questions that gauge the potential for immediate danger to the patient or others as a result of the violent incident (Table 1). The questions do not have to be used verbatim; physicians or other health care providers should choose language that is most comfortable for them but also understandably direct in gathering the perceived intentions of the patient, family, and friends. Much of this information can be asked

while obtaining the circumstances that led to the ED visit, naturally fitting into the “event” portion of the medical history. Asking first if the patient knows the other person or persons who were involved in the incident ascertains the risk for retaliation. If this is a truly random act such as mugging or robbery, then it is unlikely that the involved parties will meet again; however, such encounters are the exception rather than the rule.²⁷ Importantly, there may be some confusion around the term “know the person,” such that the youth may, in fact, recognize the assailant from the neighborhood or school but not be familiar with his or her family, friends, or “business.” Therefore, a negative response should be confirmed by asking if there is a possibility of even just meeting or seeing the other involved person(s) again. Stated intent to retaliate while in the ED conveys high risk of actual violent interactions with the involved parties in the 2 months subsequent to the ED visit, including a 5-fold risk of the youth threatening to hurt someone.²⁸ Copeland-Linder and colleagues²⁹ have shown that retaliatory attitudes ascertained during an interview soon after a violent event were associated with higher levels of aggression and fighting behavior more than 6 months after that event. Asking about whether the police or other authorities are or will be involved gauges the “protective” factor of the youth and family “delegating justice” rather than taking it into their own hands. Unfortunately, there is often distrust between the criminal justice system and poor, urban community members, in addition to a culture of “no snitching” in some areas, so the suggestion to “allow the police to handle it” may meet with some skepticism or resistance.

Assessment of Long-Term Psychosocial Risk and Protective Factors

The second component of the assessment of assault-injured youth involves multiple socioecologic domains. Given the prevalence of violence among our youth, it is crucial that we work toward programs that can reduce the immediate and long-term effects of these events. Much work has been done to identify particular risk factors and protective factors for violent injury; understanding these factors will allow for more targeted interventions. These include individual, family, and peer factors such as psychological functioning, risk, and protective behaviors including gang involvement and drug and/or alcohol use, symptoms of psychological trauma, and the adolescent’s interests, hobbies, and coping skills. The interaction between early childhood experiences and mental health is

TABLE 1. The Children’s Hospital of Philadelphia violence screening tool.

Do you know the person that hurt you?
Do you think that the fight or argument that caused this is over?
Do you plan to hurt anyone because of what happened?
Do you think that your friends or family will hurt anyone because of what happened?
Have you reported this incident to the police or other authority, and if not, do you want to?

undeniable, and this blurs the line between psychosocial and environmental risk factors.³⁰⁻³³ Asking such personal questions of an adolescent with whom we do not have an existing relationship takes skill and understanding. Certainly, adolescents, particularly those youth who have been or are currently involved in the criminal or juvenile justice systems, are cautious about revealing themselves to unfamiliar adults. The person chosen in the hospital setting to conduct screening and assessments and make referrals to outside agencies must be seen as trustworthy and nonjudgmental. Moreover, the assessments should not be “done” to the patients and their families; rather, the adolescents and their families must be actively engaged as partners in figuring out how to prevent any future violent events or stressful life experiences.²⁰ To assist in this task, several violence-related screening and assessment instruments have been successfully used in the hospital setting, some of which can be administered via a computer, and cover multiple domains associated with assaultive injury including but not limited to the following: retaliation, peer aggression, attitudes toward fighting and aggression, alcohol and substance abuse, psychological trauma, and internalizing and externalizing behaviors including depression, parental involvement, and future orientation.^{10,25,26,34-37}

Demographic Risk Factors

Multiple demographic risk factors including sex, race, family composition, educational level of the youth and caretakers, and income have all been associated with the risk of violent injury. For example, boys are more often involved in community violence outside dating relationships.³⁸ Of note, the influence of environmental and behavioral factors may render some demographic variables less important in predicting violent behavior.³⁹ For example, although black race is often cited as having a high correlation with violent injury, investigators have shown that this becomes less important after adjustment for income, sex, perception of intent, and exposure to violence.^{38,40} Similarly, high mobility (moved primary residence 2 times in the past 4 years) may be associated with a disrupted family, which can also place the child at risk for perpetration of and victimization by violence,^{38,41} as can family dysfunction, family violence, and inadequate parental monitoring.⁴² Some demographic factors such as access to a gun in the home are more clearly associated with violent injury even after other risk indicators are statistically controlled.^{43,44} Certainly, living in a neighborhood

with high rates of crime and a high level of concentrated poverty increases the risk of adolescent injury and violent perpetration.⁴⁵ Exposure to violence on television and other media may reinforce false stereotypes,⁴⁶ which is particularly problematic for youth who have been repeatedly told they cannot succeed and feel that the only way they can secure basic positive status, or “juice,” is through violence.⁴⁷ Moreover, violence is often perceived by youth in the inner city as a way to prevent future victimization.¹⁸

Psychosocial Characteristics

An individual's attitude or temperament can impact the risk of violent injury. Assessment of the circumstances under which the youth would engage in violence can be very important in determining that child's overall risk.^{38,48} For example, asking “what situation would make you fight” is a way of assessing the child's reactivity. Answers to this question, such as “if someone looks at me the wrong way,” are more concerning than “if someone hits me first.” Similarly, asking about grades, how often he or she misses school, and how they feel about teachers or other students at school are also associated with violence risk.^{41,49,50} School connectedness or school engagement including being close to others at school, feeling like part of the school, and feeling that teachers treat students fairly is perhaps the most important protective factor in minimizing the risk of subsequent involvement in violent activities.⁵¹ If there are significant issues that relate to a particular school, most urban centers have school advocacy programs that can assist students and their parents/guardians in finding and enrolling in the most appropriate school. Some, but certainly not all, school districts want to avoid recurrences of violence and can be helpful in identifying the best school setting. Moreover, school disengagement or dropping out of school is powerfully associated with delinquent and antisocial behavior.^{41,48} Violent youth, especially girls, are also more likely to have poor mental health compared with their nonviolent counterparts.⁴⁸ Self-reported depression, anxiety, and stress are associated with handgun carrying, which, in turn, significantly increases the youth's risk of violent injury.⁴⁰

Behavioral Characteristics

The behavioral risk factor most strongly associated with youth violence is weapon carrying. DuRant and colleagues⁵² have shown that carrying a weapon is a very strong indicator of frequency of fighting and

a tendency to use violence to resolve interpersonal conflict. Furthermore, there is considerable evidence that weapon carrying among teenagers is quite prevalent. In a study of more than 12 000 students in grades 7 through 12, more than 12% report carrying a weapon to school during the month before the interview.⁵⁰ In another study, 42% of urban teenagers said that they could acquire a gun if they wanted one, and 17% had carried a concealed gun.⁵³ Many youths carry a weapon to feel safer in their environment and may hope never to need to use it. Nevertheless, delinquent or deviant behavior, most often defined as a history of theft, destruction of property, or suspension or expulsion from school, is highly correlated with violent injury,⁵⁰ and youth with injuries from violence that result in ED visits report high rates of past-year peer violence and assault-related injury.⁵⁴ Youths who report selling illicit drugs are also at very high risk for being injured violently.⁵⁰ Persistently violent youth are 10 times as likely to sell drugs as are nonviolent youth.⁴⁸

Other problem behaviors of adolescence may offer a clue about the youth likely to experience violence. Alcohol use, both acute and chronic, is more commonly reported in patients with violent injuries compared with those with nonviolent injuries. Acute or binge drinking was found to be of more consequence than chronic use.⁵⁵ Tobacco or drug use, including marijuana or cocaine, has been found to correlate highly with youth's intentions to use violence in various situations and with ED visits for assault-related injuries.^{54,56} The association between drug use and violence is impressive: students who reported using tobacco or marijuana in the 9th grade were more likely to carry a gun by the time they reached the 12th grade.⁴⁰

Environmental Characteristics

Environmental conditions that place a youth at risk for violence relate to past or recent exposure to violence and community norms regarding violent behavior.³⁹ Although the nature of the term "environmental" suggests that many of these conditions can be altered, a lifetime of negative experiences is more difficult to counteract. Prior victimization or injury caused by community violence strongly predicts future perpetration and victimization.^{38,50} Witnessing violence in the community has similar effects; one may even consider a dose-response relationship in this regard. For example, witnessing an actual knife or gun attack is more influential in predicting male violent behavior than are other types of exposure such as

witnessing school or home violence.³⁹ This does not necessarily mean that the exposure actually causes subsequent violent behavior, but may indicate that the youth exists in a community where this type of event can occur as part of daily routine. Some authors propose a contagion model of violence, whereby increasing threats and exposure to violence causes more youth to carry weapons, which, in turn, further increases the perceived threat within the community, creating a vicious cycle.⁵³ Violence exposure through the media can also have deleterious effects on our youth. It has been shown that children who watch 6 or more hours of television are more likely to exhibit symptoms of depression, anxiety, and violent behaviors.⁵⁷

Experiencing or witnessing family violence shows a smaller but still significant contribution to a youth's risk for community and family violence.^{38,39} Family structure and integrity also play a role, in that the presence of both parents in the household is protective rather than predictive of violent injury.^{38,53} The level of affection and support from parents and peers are also important.⁴¹ Family functioning is also important to assess because family cohesion is a known protective factor and family violence a known risk factor. Therefore, it is important to obtain information about negative life events such as separation or divorce, remarriage, recent family suicide attempt or completion, death of a family member, and parental loss of a job.^{41,50}

INTERVENTIONS, OPPORTUNITIES, AND STRATEGIES

Family engagement is crucial to the success of most interventions; however, often times, a parent does not know how to satisfactorily manage, support, and monitor his/her child, and many parents are going to need help in keeping their son/daughter out of harm's way. The ED team can support the family by providing brief interventions after initial assessment, giving appropriate postdischarge contacts and connections, and engaging the patient and family into community-based programs.

Brief Intervention in the ED

Effective brief interventions, generally based on the principles and practices of motivational interviewing, are available for multiple problem areas and have been successfully used in the hospital setting.⁵⁸ Cunningham and colleagues⁵⁹ demonstrated the effectiveness of a single therapist conducted brief intervention session at a level I trauma center in reducing peer victimization and

aggression and adverse consequences from alcohol use among adolescents who had engaged in some form of violence during the prior year. Using a randomized control trial design, some of these differences held up at 12 months postdischarge. Similarly, Johnston and colleagues⁶⁰ used a single behavior change counseling session with adolescent patients who were primarily involved in unintentional injuries. They found that patients who received such sessions were more likely to use seat belts and wear bicycle helmets than a randomly assigned comparison group. Although these studies are encouraging, the basic lesson learned is that young people can make some, although limited, behavioral changes as a result of a brief intervention in the hospital setting after medical treatment. Not surprisingly, programs that engage violently injured youth in the hospital setting, conduct screenings and assessments, and refer patients to ongoing wraparound and case management services for 6 to 12 months after hospital release have secured more robust changes and outcomes regarding subsequent victimization and violent perpetration.⁶¹⁻⁶³ Of course, it is optimal to link patients and their families to community-based programs if they are available, as follow-up on the initial work done at the hospital.³⁵ This requires “strategic collaboration” and the adoption of a thoroughgoing public health approach to violence prevention.²⁰

Community-Based Interventions

Interventions based on strategies such as scare tactics, boot camps, gun buy-backs, and isolated self-esteem enhancement programs that provide information without skills were ineffective or even harmful.^{64,65} Although well-meaning hospitals and medical providers would like teens to understand “what might happen,” research shows that programs for youth that tour trauma bays and morgues are not effective prevention strategies and may actually cause more harm than good by retraumatizing the participants. The most successful programs evaluated were based on strategies such as social skills training, positive youth development, mentoring, parent and family training, and home visitation. Effective violence prevention approaches, short of massive social and economic reform, build resilience and enhance protective factors to overcome social and environmental stressors.⁶⁶⁻⁶⁹ These approaches include encouraging participation in prosocial peer groups, appropriate school settings, and community-based programs that emphasize positive social norms; facilitating concomitant involvement in safe activities; providing supportive

relationships with positive adults in their communities; and enhancing competence in cognitive, social, and emotional skills.

Family-focused interventions such as multisystemic therapy and functional family therapy are among the most effective interventions for chronically violent youth.^{70,71} This is not to suggest that these kinds of programs need to be provided at the hospital, but rather, to ensure that the hospital's social work team is aware of such programs in the community. Both programs focus centrally on family dynamics, both use cognitive behavioral techniques, both programs use therapists who work with the adolescents in their own neighborhoods, and both programs work diligently to engage adolescents in school and other prosocial activities. One promising violence reduction program—Cease-Fire Chicago, now known as Cure Violence—has been adopted in at least one major hospital-based program.⁷² Cure Violence is implemented in high-crime neighborhoods and uses outreach workers and interventionists who work with individuals known to be at highest risk for perpetrating gun violence. The outreach workers and interventions are chosen based on their “street credibility” and their skills and passion for reducing street violence. Multisystemic therapy and functional family therapy, as well as Cure Violence, include elements common to many hospital-based violence intervention programs, particularly the core model Caught in the Crossfire, that have also been shown to reduce retaliatory violence and re-injury.

In the prevention literature, meta-analyses typically focus on programs or risk factors as the primary unit of analysis. A recent meta-analysis used program components as the unit of analysis to detect which components are most effective among youth violence prevention programs in reducing violence recidivism.⁷³ Program characteristics that discern which programs are likely to be most successful include focus, intensity, quality, fidelity, emphasis on cognitive behavioral approaches, appropriate response to risk level, and the use of a written procedures manual. Although this protocol needs more refinement, the identified elements are feasible to adopt via hospital-based violence intervention programs or hospital-affiliated programs that provide or refer patients to needed services and case management. These programs identify, assess, and support assault-injured patients after discharge from the hospital.⁷⁴ Many of these programs have joined to form a coalition, The National Network of Hospital-based Violence Intervention Programs (www.nnhvip.org). These programs share data and best practices in administration and

community-level intervention and seek to recruit and help build similar programs across the country. The network has developed a “best practices guide” that outlines the steps required to start and maintain this kind of program within a hospital system.⁷⁵

SUMMARY

The visit by an assault-injured youth in the emergency department setting is an opportunity to break the immediate and potentially long-term consequences of violence. Individual emergency medical practitioners can have a substantial impact on the way that these patients respond to medical and psychosocial evaluation. Hospital systems can support their patients and the communities in which they exist by designing and implementing trauma-informed programs that fully assess, refer, and even case manage these youth after hospital discharge. ❏

REFERENCES

- Klaus P, Rennison C. Age patterns in violent victimization, 1976-2000. Washington, DC: United States Department of Justice, Bureau of Justice Statistics; 2002. Available at: <http://bjs.ojp.usdoj.gov/content/pub/pdf/apvv00.pdf>. Accessed 1/15/13.
- Kennedy D. Don't shoot: one man, a street fellowship, and the end of violence in inner-city America. New York, NY: Bloomsbury; 2011.
- Smith D, Ecob R. An investigation into causal links between victimization and offending in adolescents. *Brit J Sociol* 2007; 58:633-59.
- Chang JJ, Chen JJ, Brownson RC. The role of repeat victimization in adolescent delinquent behaviors and recidivism. *J Adolesc Health* 2003;32:272-80.
- Jennings WG, Higgins GE, Tewksbury R, et al. A longitudinal assessment of the victim-offender overlap. *J Interpers Violence* 2010;25:2147-74.
- Rich J. Wrong place, wrong time: trauma and violence in the lives of young black men. Baltimore, MD: John Hopkins University Press; 2009.
- Task Force on Adolescent Assault Victim Needs. Adolescent assault victim needs: a review of issues and a model protocol. *Pediatrics* 1996;98:991-1000.
- Muelleman RL, Reuwer J, Sanson TG, et al. An emergency medicine approach to violence throughout the life cycle. SAEM Public Health and Education Committee. *Acad Emerg Med* 1996;3:708-15.
- Johnson SB, Bradshaw CP, Wright JL, et al. Characterizing the teachable moment: is an emergency department visit a teachable moment for intervention among assault-injured youth and their parents? *Pediatr Emerg Care* 2007;23:553-9.
- Cunningham R, Knox L, Fein J, et al. Before and after the trauma bay: the prevention of violent injury among youth. *Ann Emerg Med* 2009;53:490-500.
- Centers for Disease Control and Prevention. 10 Leading causes of death and injury. Available at: <http://www.cdc.gov/injury/wisqars/LeadingCauses.html>. Accessed November 2, 2012.
- Centers for Disease Control and Prevention. Nonfatal injury data. Available at: <http://www.cdc.gov/injury/wisqars/nonfatal.html>. Accessed November 2, 2012.
- Cheng TL, Wright J, Fields CB, et al. Violent injuries among adolescents: declining morbidity and mortality in an urban population. *Ann Emerg Med* 2001;37:292-300.
- Campbell C, Schwartz D. Prevalence and impact of exposure to interpersonal violence among suburban and urban middle school students. *Pediatrics* 1996;98:396-402.
- Briggs-Gowan MJ, Ford JD, Fraleigh L, et al. Prevalence of exposure to potentially traumatic events in a healthy birth cohort of very young children in the northeastern United States. *J Trauma Stress* 2010;23:725-33.
- Johnson AO, Harun N, Moore CG, et al. Violence and drug use in rural teens: national prevalence estimates from the 2003 Youth Risk Behavior Survey. *J Sch Health* 2008;78:554-61.
- Berton MW, Stabb SD. Exposure to violence and post-traumatic stress disorder in urban adolescents. *Adolescence* 1996;31:489-98.
- Rich JA, Stone DA. The experience of violent injury for young African-American men: the meaning of being a “sucker”. *J Gen Int Med* 1996;11:77-82.
- Dowd DM, Seidel JS, Sheehan K, et al. Teenagers' perceptions of personal safety and the role of the emergency health care provider. *Ann Emerg Med* 2000;34:346-50.
- Denninghoff KR, Know L, Cunningham R, et al. Emergency medicine: competencies for youth violence prevention and control. *Acad Emerg Med* 2002;9:947-56.
- Rice C. Power concedes nothing. New York, NY: Scribner; 2012.
- Rich J, Corbin T, Bloom S, et al. Healing the hurt: trauma informed approaches to the health of boys and young men of color. California Endowment. Available at: <http://www.calendow.org/uploadedFiles/Publications/BMOC/Drexel%20-%20Healing%20the%20Hurt%20-%20Full%20Report.pdf>. Accessed November 21, 2011.
- Chamberlain LB. The amazing adolescent brain: trauma and the potential for healing. Available at: http://www.instituteforsafefamilies.org/pdf/theamazingbrain/The_Amazing_Brain-2.pdf. Accessed November 21, 2011.
- Bloom SL, Bennington-Davis M, Farragher B, et al. Multiple opportunities for creating sanctuary. *Psychiatr Q* 2003;74: 173-90.
- Fein JA, Kassam-Adams N, Gavin M, et al. Persistence of posttraumatic stress in violently injured youth seen in the emergency department. *Arch Pediatr Adolesc Med* 2002;156: 836-40.
- Fein JA, Kassam-Adams N, Vu T, et al. Emergency department evaluation of acute stress disorder symptoms in violently injured youth. *Ann Emerg Med* 2001;38:391-6.
- Cheng TL, Johnson S, Wright JL, et al. Assault-injured adolescents presenting to the emergency department: causes and circumstances. *Acad Emerg Med* 2006;13(6):610-6.
- Wiebe DJ, Blackstone MM, Mollen CJ, et al. Self-reported violence-related outcomes for adolescents within eight weeks of emergency department treatment for assault injury. *J Adolesc Health* 2011;49:440-2.
- Copeland-Linder N, Johnson SB, Haynie DL, et al. Retaliatory attitudes and violent behaviors among assault-injured youth. *J Adolesc Health* 2012;50:215-20.
- Dube SR, Anda RF, Felitti VJ, et al. Childhood abuse, household dysfunction, and the risk of attempted suicide throughout the life span: findings from the Adverse Childhood Experiences Study. *JAMA* 2001;286:3089-96.

31. Dube SR, Felitti VJ, Dong M, et al. Childhood abuse, neglect, and household dysfunction and the risk of illicit drug use: the adverse childhood experiences study. *Pediatrics* 2003;111:564-72.
32. Anda RF, Brown DW, Felitti VJ, et al. Adverse childhood experiences and prescribed psychotropic medications in adults. *Am J Prev Med* 2007;32:389-94.
33. Anda RF, Felitti VJ, Bremner JD, et al. The enduring effects of abuse and related adverse experiences in childhood. A convergence of evidence from neurobiology and epidemiology. *Eur Arch Psychiatry Clin Neurosci* 2006;256:174-86.
34. Mollen CJ, Fein JA, Localio AR, et al. Characterization of interpersonal violence events involving young adolescent girls vs events involving young adolescent boys. *Arch Pediatr Adolesc Med* 2004;158:545-50.
35. Aboutanos MB, Jordan A, Cohen R, et al. Brief violence interventions with community case management services are effective for high-risk trauma patients. *J Trauma* 2011;71:228-36.
36. Anixt JS, Copeland-Linder N, Haynie D, et al. Burden of unmet mental health needs in assault-injured youths presenting to the emergency department. *Acad Pediatr* 2012;12:125-30.
37. Redeker N, Smeltzer S, Kirkpatrick J, et al. Risk factors of adolescent and young adult trauma victims. *Am J Crit Care* 1995;4:370-8.
38. Malik S, Sorenson SB, Aneshensel CS. Community and dating violence among adolescents: perpetration and victimization. *J Adolesc Health* 1997;21:291-302.
39. Song LY, Singer MI, Anglin TM. Violence exposure and emotional trauma as contributors to adolescents' violent behaviors. *Arch Pediatr Adolesc Med* 1998;152:531-6.
40. Simon TR, Richardson JL, Dent CW, et al. Prospective psychosocial, interpersonal, and behavioral predictors of handgun carrying among adolescents. *Am J Public Health* 1998;88:960-3.
41. Saner H, Ellickson P. Concurrent risk factors for adolescent violence. *J Adolesc Health* 1996;19:94-103.
42. McCord J, Widom C, Crowell N, editors. *Juvenile crime, juvenile justice*. Washington, DC: National Academy Press; 2001. p. 167-76.
43. Ruback R, Shaffer J, Clark VA. Easy access to firearms: juveniles' risks for violent offending and violent victimization. *J Interpers Violence* 2011;26:2111-38.
44. Kellerman AL, Rivara FP, Rushforth N, et al. Gun ownership as a risk factor for homicide in the home. *N Engl J Med* 1993;329:1084-91.
45. MacDonald J, Gover AR. Concentrated disadvantage and youth-on-youth homicide: assessing the structural covariates over time. *Homicide Studies* 2005;9:30-54.
46. Strasburger VC, Donnerstein E. Children, adolescents, and the media: issues and solutions. *Pediatrics* 1999;103:129-39.
47. Johnson SB, Frattaroli S, Wright JL, et al. Urban youths' perspectives on violence and the necessity of fighting. *Inj Prev* 2004;10:287-91.
48. Ellickson P, Saner H, McGuigan KA. Profiles of violent youth: substance use and other concurrent problems. *Am J Public Health* 1997;87:985-91.
49. Loeber R, Farrington DP. Young children who commit crime: epidemiology, developmental origins, risk factors, early interventions, and policy implications. *Dev Psychopathol* 2000;12:737-62.
50. Resnick MD, Bearman PS, Blum RW, et al. Protecting adolescents from harm: findings from the National Longitudinal Study on Adolescent Health. *JAMA* 1997;278:823-32.
51. Lösel F, Farrington DP. Direct protective and buffering protective factors in the development of youth violence. *Am J Prev Med* 2012;43S8-23.
52. DuRant RH, Kahn J, Beckford PH, et al. The association of weapon carrying and fighting on school property and other health risk and problem behaviors among high school students. *Arch Pediatr Adolesc Med* 1997;151:360-6.
53. Bergstein J, Hemenway D, Kennedy B, et al. Guns in young hands: a survey of urban teenagers' attitudes and behaviors related to handgun violence. *J Trauma Inj Infect Crit Care* 1996;41:794-8.
54. Ranney M, Whiteside L, Walton M, et al. Sex differences in characteristics of adolescents presenting to the emergency department with acute assault-related injury. *Acad Emerg Med* 2011;18:1027-35.
55. Borges G, Cherpitel CJ, Rosovsky H. Male drinking and violence-related injury in the emergency room. *Addiction* 1998;93:103-12.
56. DuRant RH, Treiber F, Goodman E, et al. Intentions to use violence among young adolescents. *Pediatrics* 1996;98:1104-8.
57. Singer MI, Slovak K, Frierson T, et al. Viewing preferences, symptoms of psychological trauma, and violent behaviors among children who watch television. *J Am Acad Child Adolesc Psychiatry* 1998;37:1041-8.
58. Burke BL, Arkowitz H, Menchola M. The efficacy of motivational interviewing: a meta-analysis of controlled clinical trials. *J Consult Clin Psychol* 2003;71:843-61.
59. Cunningham RM, Chermack ST, Zimmerman MA, et al. Brief motivational interviewing intervention for peer violence and alcohol use in teens: one-year follow-up. *Pediatrics* 2012;129:1083-90.
60. Johnston BD, Rivara FP, Droesch RM, et al. Behavior change counseling in the emergency department to reduce injury risk: a randomized, controlled trial. *Pediatrics* 2002;110:267-74.
61. Shibu D, Zahnd E, Becker M, et al. Benefits of a hospital-based peer intervention program for violently injured youth. *J Am Coll Surg* 2007;205:684-9.
62. Cooper C, Eslinger DM, Stolley PD. Hospital-based violence intervention programs work. *J Trauma* 2006;61:534-40.
63. Zun LS, Downey L, Rosen J. The effectiveness of an ED-based violence prevention program. *Am J Emerg Med* 2006;24:8-13.
64. Commission for the Prevention of Youth Violence. *Youth and violence. Medicine, nursing, and public health: connecting the dots to prevent violence*. Available at: <http://www.scribd.com/doc/50914270/Connecting-the-Dots-to-Prevent-Youth-Violence>. Accessed November 8, 2012.
65. U.S. Department of Health and Human Services. *Youth violence: a report of the surgeon general*. Rockville, MD: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Injury Prevention and Control; 2001.
66. Catalano RF, Loeber R, McKinney KC. *School and community interventions to prevent serious and violent offending*. Washington, DC: US Department of Justice, Office of Justice Programs, Office of Juvenile Justice and Delinquency Prevention; 1999.
67. Catalano RF, Arthur MW, Hawkins JD, et al. Comprehensive community- and school-based interventions to prevent antisocial behavior. In: Loeber R, Farrington DP, editors. *Serious and violent juvenile offenders: risk factors and successful interventions*. Thousand Oaks, CA: Sage Publications; 1998. p. 248-83.
68. Masten AS, Coatsworth JD. The development of competence in favorable and unfavorable environments: lessons from research on successful children. *Am Psychol* 1998;53:205-20.

69. Catalano RF, Hawkins JD. *Communities that care: risk-focused prevention using the social development strategy*. Seattle, WA: Developmental Research and Programs, Inc.; 1995.
70. Henggeler SW, Clingempeel WG, Brondino MJ, et al. Four-year follow-up of multisystemic therapy with substance-abusing and substance-dependent juvenile offenders. *J Am Acad Child Adolesc Psychiatry* 2002;41:868-74.
71. Sexton TL, Alexander JF. *Functional family therapy*. Washington, DC: Juvenile Justice Bulletin; 2000; Available at: <https://www.ncjrs.gov/pdffiles1/ojdp/184743.pdf>. Accessed 1/15/13.
72. Webster DW, Whitehill JM, Vernick JS, et al. Effects of Baltimore's safe streets program on gun violence: a replication of Chicago's ceasefire program. *J Urban Health* 2013;90:27-40.
73. Lipsey M. The primary factors that characterize effective interventions with juvenile offenders: a meta-analytic overview. *Victims Offenders* 2009;4:124-47.
74. Snider C, Lee J. Youth violence secondary prevention initiatives in emergency departments: a systematic review. *Can J Emerg Med* 2009;11:161-8.
75. Karraker N, Cunningham RK, Becker MG, et al, editors. *Violence is preventable: a best practices guide for launching and sustaining a hospital-based program to break the cycle of violence*. Washington, DC: Office of Victims of Crime, Office of Justice Programs, U.S. Department of Justice; 2011, Available at: <http://youthalive1.squarespace.com/storage/ViolenceisPreventableOnline.pdf>. Accessed 1/15/13.